

Combined Effect of Topology and Synchronism Perturbation on Cellular Automata: Preliminary Results

Jean-Baptiste Rouquier¹ and Michel Morvan^{1,2}

¹ Université de Lyon, IXXI and ENS Lyon, France

² EHESS and Santa Fe Institute.

Abstract. The aim of this paper is to experimentally study the combined effect of the introduction of two kinds of structural perturbations to the behavior of cellular automata. We present the results obtained by simultaneously perturbing synchronism and topology of elementary cellular automata. We show that very interesting and different behaviors appear, including phase transitions and non monotonicity (i.e. introduction of both perturbations is less effective than the introduction of only one of them). These results lead us to think that this study is worth to be now developed more accurately.

1 Introduction

The last decade has seen the development of an interest to the effect of structural perturbation applied to discrete dynamical model [1–5]. Almost all such studies were focused on the introduction of one kind of perturbation. However, in [6], it is shown that combining two perturbations, namely the introduction of asynchronism and a topology perturbation, could lead to interesting new behaviors of the Game of Life. In particular, it has been shown that while the Game of Life was very sensitive to asynchronism, this sensitivity was strongly decreased when coupled with the breaking of a small percentage of links in the grid.

The aim of this paper is to develop an exhaustive study of the coupling of the same kind of perturbations for elementary cellular automata. We will show that very different behaviors appear. On the one hand, part of these results were expected: some ECA are robust to both perturbations, even when applied simultaneously, other are sensitive to only one, the other having no effect, and some are sensitive to both of them when they are applied simultaneously. On the other hand, some interesting and more unexpected behavior appear in this last case. For example, the combination of both perturbation can in some cases lead either to the vanishing of a phase transition induced by one perturbation alone or to the appearance of a new phase transition. Moreover, rule 45 exhibits the same behavior as the Game of Life: one perturbation makes the system robust

This work has been partially supported by French ANR project NT05-2-4913 CARPVIRTUEL and European NEST Pathfinder project 043322 MORPHEX.

to the introduction to the other, and so the effects of the perturbations are not combined in a monotonous way.

The main conclusion of this exploratory paper is that the robustness to the combination of different perturbations is a subject which deserves to be studied in more details, even in very simple systems as ECA.

The paper is organized as follows. In Section 2, we describe the definitions and in particular the definition of the perturbations we will apply to ECA. We also present the experimental protocol. Section 3 contains the result and brief analysis of the experimental results. We classify the different observed behavior and highlight those that are, in our opinion, the most interesting for future studies. We end by a discussion in which we propose promising research directions.

2 Definitions and Experimental Protocol

2.1 Perturbed Cellular Automata

Definition 1. A Cellular Automata (CA) with periodic boundary condition is a tuple consisting in

- Q a set of states,
- $\mathcal{U} := (\mathbb{Z}/n\mathbb{Z})^d$ the finite set of cells (d is the dimension, n is a parameter),
- $V := \{\vec{v}_1, \dots, \vec{v}_{|V|}\}$ is the neighborhood, a finite set of vectors of \mathbb{Z}^d ,
- $\delta : Q^{|V|} \rightarrow Q$ is the update rule.

A *configuration* specifies the state of each cell, and is thus a function $\mathcal{U} \rightarrow Q$. The usual dynamics on a CA is the following: considering a configuration c_t , the next configuration c_{t+1} is defined by $\forall z, c_{t+1}(z) := \delta(c_t(z + \vec{v}_1), \dots, c_t(z + \vec{v}_{|V|}))$.

This early works focuses on a particular class of CA, namely the Elementary Cellular Automata (ECA), the simplest ones: $Q = \{0, 1\}$, $d = 1$, $V = \{-1, 0, 1\}$. We use the classical notation introduced by Wolfram: an ECA δ is denoted by the code $\delta(0, 0, 0) \times 2^0 + \delta(0, 0, 1) \times 2^1 + \delta(0, 1, 0) \times 2^2 + \dots + \delta(1, 1, 1) \times 2^7$.

The fact that it is one dimensional implies that n is also the number of cells.

We perturb this dynamics in two ways. First, not all cells are updated at each time step. Instead, at each time step, each cell has a fixed probability p to get updated, independently of the other cells. p is a parameter. The case $p = 1$ is the unperturbed (synchronous) case. Updating only one cell at a time would also be a classical model [7].

Second, a cell might not always “see” all its neighbors. Instead, at each time step, when the cell z requests the states of its neighbors to compute its next state, it gets no answer with probability r . It then assumes the neighbor is in state q . q and r are parameters. The case $r = 0$ (no link cut) is the unperturbed case.

The final dynamics is thus:

$$c_{t+1} := \begin{cases} \delta(c'_t(z-1), c_t(z), c'_t(z+1)) & \text{with probability } p \\ c_t(z) & \text{with probability } 1 - p \end{cases} \quad (1)$$

where $c'_t(y) := \begin{cases} q & \text{with probability } r \\ c_t(y) & \text{with probability } 1 - r \end{cases}$

Motivation of the Model Not updating a cell can be seen as a delay before updating. It can also be seen as a defective cell, where we chose that a defective cell keeps its state. When a cell does not sees one of its neighbor, it corresponds to a defective (oriented) link between the cells.

All defects are temporary. This is necessary for ECA since otherwise a defect would prevent all communication between its left and right sides.

2.2 Experimental Protocol

We call *run* the temporal evolution of a CA when all parameters are chosen. The macroscopic measure we use on a run is the density of cells in state 1. This measure has been shown relevant in [8]. For a configuration c , we note the density

$$\rho(c) := \#\{z \in \mathcal{U} \mid c(z) = 1\} / |\mathcal{U}|$$

Sampled Parameters We do one run for each possible combination of the following parameters.

For a given choice of the rule (including the parameter q), we sample many values of p and r , that is, we combine many strengths of synchronism and topology perturbation. For the former, we sample the whole space of synchrony rates, including the synchronous case, by steps of 0.01: $p \in \{0.01, 0.02, \dots, 1.00\}$. For the latter, cutting 10% of the links is already a strong perturbation, and we do not perturb the model more. That is, $r \in \{0.00, 0.01, 0.02, \dots, 0.10\}$.

We sample both values (0 and 1) for the parameter q . For a given rule, e.g. 110, we note the parameter q as a subscript, e.g. 110_0 and 110_1 . For autoconjugate rules (i.e. rules that are unchanged under the exchange of state 0 and 1), both choices of the parameter q leads to the same dynamics (actually to symmetric ones). We therefore study only one, which we note 105_a .

When working on ECA, one usually considers only 88 rules among the 256 possible ones, thanks to symmetry considerations. The left/right symmetric of a rule δ (i.e. the rule $\delta'(a, b, c) := \delta(c, b, a)$) leads to configurations symmetric to the ones obtained with δ . Therefore, we will only consider one rule among each pair (δ, δ') . However, the other usual symmetry, namely exchanging the states 0 and 1, is no longer possible. Indeed, the parameter q has introduced an asymmetry between both states.

Fixed Parameters The following parameters are identical for all runs.

- The ring size is $n=10\,000$ cells. [4] estimated that 2 000 was already sufficient in a similar problem.
- The initial configuration is random (each cell is in state 0 with probability 0.5, independently from the other cells) and distinct for each run.
- To speed up computation, we do not measure ρ at each time step, instead we do the same number of measurements (roughly 200) in each decade.

- To detect when the density has settled to a constant value, we do a linear regression on the last 200 measure points and wait for the fitted line to be almost horizontal (we actually do three independent fits on three portions of the history). We then average the density over the history.
- If the density has not stabilized after 10^6 steps, we stop the simulation and ignore this point.

Limits of the Protocol The density might be seen as a quite rough measure, and so we will certainly miss some phenomenons occurring in this model. However, the variety of results shows that this measure is able to detect one kind of sensibility to the perturbations applied here.

Another limit is that this protocol does not measure the true asymptotic density, but instead a long enough stage where the density is stable. This is the only stage observable experimentally if the asymptotic regime occur only after exponential time. Nevertheless, decision to stop a run might occur early if the density is very slowly decreasing.

Last, for each value of the parameters, we compute only one data point. The smoothness of most surfaces shows that variance among runs is low, and that we do not need to average over several data points.

3 Exhaustive Study of the ECA Space

We now apply this protocol to each ECA and plot ρ against p and r (Fig. 1). Reviewing the sampling surfaces suggests the classification in four classes of Table 1, based on the sensitivity to each perturbation. The subclasses a/, b/ and c/ are explained in Section 3.4. We now review each class.

		Topology perturbation																					
		insensitive								sensitive													
Synchronism perturbation insensitive		0 ₀	0 ₁	2 ₀	8 ₀	8 ₁	10 ₀	23 _a	4 ₁	5 ₀	12 ₁	13 ₀	15 _a	29 ₀	29 ₁	32 ₁	36 ₁	40 ₁	44 ₁	72 ₀	72 ₁		
		24 ₀	28 ₀	32 ₀	34 ₀	40 ₀	42 ₀		76 ₁	78 ₁	104 ₀	104 ₁	132 ₁	140 ₁	160 ₁	164 ₁	168 ₁	170 _a					
		51 _a	56 ₀	60 ₀	60 ₁	74 ₀	105 _a		172 ₁	184 ₀	184 ₁	200 ₀	232 _a										
		128 ₀	128 ₁	130 ₀	136 ₀	136 ₁																	
		138 ₀	152 ₀	154 ₁	160 ₀	162 ₀																	
		168 ₀	200 ₁	204 _a																			
Synchronism sensitive		1 ₀	4 ₀	5 ₁	6 ₀	9 ₀	11 ₀	11 ₁	12 ₀	a/	2 ₁	6 ₁	10 ₁	24 ₁	34 ₁	38 ₁	42 ₁	45 ₀	45 ₁	46 ₁	56 ₁	73 ₀	73 ₁
		18 ₀	19 ₀	19 ₁	22 ₀	26 ₀	33 ₀				74 ₁	77 _a	94 ₀	94 ₁	130 ₁	134 ₁	138 ₁	152 ₁	154 ₀	162 ₁			
		35 ₀	35 ₁	36 ₀	38 ₀	41 ₀	41 ₁			b/	18 ₁	26 ₁	50 ₁	58 ₁	106 ₀	106 ₁	146 ₁	178 _a					
		44 ₀	46 ₀	50 ₀	54 ₀	58 ₀	76 ₀			c/	1 ₁	3 ₀	3 ₁	7 ₀	7 ₁	9 ₁	13 ₁	14 ₀	14 ₁	22 ₁	25 ₀	25 ₁	27 ₀
		90 ₁	108 ₀	110 ₀	110 ₁	132 ₀				27 ₁	28 ₁	30 ₀	30 ₁	33 ₁	37 ₀	37 ₁	43 _a	54 ₁	57 ₀	57 ₁	62 ₀		
		134 ₀	146 ₀	164 ₀						62 ₁	78 ₀	90 ₀	108 ₁	122 ₀	122 ₁	126 ₀	126 ₁	140 ₀	142 _a				
									150 _a	156 ₀	156 ₁	172 ₀											

Table 1. Classification of the ECA from their robustness against each perturbation




Fig. 1. Some sampling surfaces (color online). Axis x (width) represents topology perturbation, y (depth) synchronism perturbation, z (height) asymptotic density. The difference between Figures 1.a and 1.f is that in the former only the unperturbed case ($p = 1, r = 0$) is non zero, whereas in the latter there is a continuous transition from non-zero to zero, with respect to both p and r .

3.1 Rules Insensitive to Both Perturbations

Most of the rules in this class have a surface that is just 0 everywhere (Fig 1.a), sometimes with an exception in the unperturbed case ($p = 1, r = 0$). A few rules (9%) outside this class also have a noticeably different behavior in the unperturbed case, but this is left for further study.

Since they are by definition insensitive to the perturbations, the other rules of this class also have a horizontal surface, (the altitude is often around $\rho = 0.5$).

3.2 Rules Sensitive Only to Synchronism Perturbation

Some rules of this class undergo a phase transition when p changes (Fig 1.b). The conditions of a conjecture by [9] are fulfilled, so this transition should belong to the universality class of directed percolation. (For an introduction to directed percolation in CA see [3], and for a detailed review of phase transitions see [10].)

It is interesting to note that the topology perturbation slightly shifts the threshold (the position of the transition) of most of those rules.

The remaining surfaces of this class are smooth, for instance like Fig 1.c. Since these rules are almost insensitive to topology perturbation, all slices along a red line (grey for a B&W version) are nearly identical.

3.3 Rules Sensitive Only to Topology Perturbation

Some rules of this class ($4_1, 5_0, 12_1, 13_0, 36_1, 44_1, 72_0, 72_1, 76_1, 78_1, 104_0, 104_1, 132_1, 140_1, 164_1, 170_a, 172_1, 184_0, 184_1, 200_0, 232_a$) are sensitive to the introduction of topology perturbation, i.e., they have a brutal change of behavior between $r = 0$ and $r > 0$ (Fig 1.d). A few rules actually have a density that depends on p , but only when $r = 0$.

The other surfaces of this class are smooth. The only phase transitions of this class are thus first order phase transitions (i.e. discontinuity of ρ), and they are all located at the introduction of topology perturbation.

3.4 Rules Sensitive to Both Perturbations

This is the most interesting class, as we will see many different behaviors. We examine specifically how both perturbations interact.

a/ Sensitive to Introduction of Perturbation As in the class 3.3, the surfaces of this subclass exhibit a gap between $r = 0$ and $r > 0$. This is a first order phase transition, and it is not the only phase transition of class 3.4.

b/ Created or Destroyed Phase Transition We have seen in section 3.2 that one of the two ways for the density to vary when p changes is a phase transition. For the ECA of Section 3.2, this phase transition was robust against topology perturbation. Here it is not the case since the phase transition is destroyed when

$r > 0$ (Fig 1.e). We looked at the configurations (not reproduced here due to lack of space) and it seems that cutting a link in a region of cells in state 0 (i.e. in the absorbing state) corresponds to reseeding an active site at this point. The configuration is thus constantly reseeded with active sites and 0^n is no more an absorbing state. This is the case for $18_1, 26_1, 50_1, 58_1, 106_1, 146_1$

The case of 178_a is the opposite: when the topology is not perturbed, this rule is autoconjugate and can't have a preferred state. In [11], without topology perturbation, it is suggested that this rule could have a phase transition of the \mathbb{Z}_2 symmetric percolation universality class when p varies. The slightest topology perturbation introduces a bias towards 0^n , which becomes the only absorbing state. The rule then satisfies the conditions of the conjecture by [9], and should thus belong to the universality class of directed percolation. Therefore, we conclude that topology perturbation has created a new phase transition.

c/ Smooth The last subclass contains the remaining surfaces, which are smooth.

Some Remarkable Rules We now would like to point out some remarkable rules of the present class (Section 3.4).

Rule 106_0 (Fig 1.f) exhibits two phase transitions, one when p varies and the other when r varies.

Rule 90_0 (Fig 1.g) is almost insensitive to a single perturbation. However, both perturbations combined induce a noticeable decrease of the density. (This study focuses on small r , but it is not hard to see that for high enough r , rule 90_0 has a vanishing density.)

Rule 122_0 (Fig 1.h) has a density close to 0.5 in the unperturbed case. When applying any one of the perturbations, the density tends to 0.6. But when applying both, the density tends to back 0.5. (However, the configurations are quite different from the unperturbed case, only the density has been restored.) One perturbation can be seen as a way to make the CA robust against the other perturbation. This is one of the rare cases where perturbations are not combining in a monotonous way.

Rule 73 (which has the same qualitative behavior for both values of q) has a constant asymptotic density when topology is not perturbed, for all values of p (except $p = 1$). With a small perturbation of the topology, ρ spans the range 0.3 to 0.5. The interesting effect is that introducing topology perturbation has made the rule sensitive to synchronism perturbation. Same remark applies to rules $14_0, 14_1, 57_0, 90_0, 150_a, 156_0, 156_1$ and to almost all rules of subclass a/.

The opposite behavior also exists: 45, with no topology perturbation, has a density that depends on p . With $r > 0$, the density becomes constant for all p : perturbing the topology makes the rule robust against synchronism perturbation.

4 Discussion and Perspectives

By means of *dynamic* perturbation, we studied topology perturbation on ECA and its interplay with synchronism perturbation. Topology perturbation is a

perturbation on the neighboring cells, while synchronism perturbation is a perturbation on the central cell. The macroscopic parameter chosen, namely the density, already allows one to see a rich set of behaviors, even if there may be more than what this measure reveals. This is both a confirmation of the relevancy of the density and a confirmation that this model could be studied further, possibly with analytical results.

There does not seem to be any relation between this classification and classical classifications about synchronous deterministic CA.

The topology perturbation introduced in this paper should be studied for itself, possibly with the following generalization of the parameter q . When a cell does not know the state of its neighbor, it currently assumes it is in state q . It could instead decide its state according to a Bernoulli law. Future work will study the combined effect of r and this new parameter q .

A few surfaces contains the characteristic curve of second order phase transition. Future work should check if they belong to the universality class of directed percolation, according to the aforementioned conjecture.

References

1. Fatès, N., Morvan, M.: An experimental study of robustness to asynchronism for elementary cellular automata. *Complex Systems* **16** (2005) 1–27
2. Ingerson, T.E., Buvel, R.L.: Structure in asynchronous cellular automata. *Physica D Nonlinear Phenomena* **10** (January 1984) 59–68
3. Fatès, N.A.: Asynchronism induces second order phase transitions in elementary cellular automata. *Journal of Cellular Automata* (march 2007)
4. Rouquier, J.B., Morvan, M.: Coalescing cellular automata: Synchronization by common random source for asynchronous updating. *Journal of Cellular Automata* (accepted, 2008)
5. Schönfisch, B., de Roos, A.: Synchronous and asynchronous updating in cellular automata. *Biosystems* **51**(3) (1999) 123–143
6. Fatès, N., Morvan, M.: Perturbing the topology of the game of life increases its robustness to asynchrony. In Sloot, P.M.A., Chopard, B., Hoekstra, A.G., eds.: 6th International Conference on Cellular Automata for Research and Industry Proceedings. Volume 3305 of LNCS., Springer (2004) 111–120
7. Fatès, N., Morvan, M., Schabanel, N., Thierry, E.: Fully asynchronous behavior of double-quiescent elementary cellular automata. *Theoretical Computer Science* **362** (2006) 1–16
8. Fatès, N.: Experimental study of elementary cellular automata dynamics using the density parameter. In: *Discrete models for complex systems, DMCS '03 (Lyon)*. Discrete Mathematics Theoretical Computer Science Proceedings, AB. Assoc. Discrete Math. Theor. Comput. Sci., Nancy (2003) 155–165
9. Grassberger, P.: Are damage spreading transitions generically in the universality class of directed percolation? *J Stat Phys* **79** (September 1995) 13–23
10. Hinrichsen, H.: Nonequilibrium critical phenomena and phase transitions into absorbing states. *Advances in Physics* **7** (Nov 2000) 815–958
11. Fatès, N.: Robustesse de la dynamique des systèmes discrets : le cas de l'asynchronisme dans les automates cellulaires. PhD thesis, ENS Lyon (December 2004)